

Digital Library

Environmental Weed Risk Assessments

Biosecurity, pests, weeds and diseases research

2022

Phalaris aquatica - environmental weed risk assessment 2022

Department of Primary Industries and Regional Development, Western Australia

Follow this and additional works at: https://library.dpird.wa.gov.au/bs_wra

Part of the Biosecurity Commons, and the Weed Science Commons

Recommended Citation

Department of Primary Industries and Regional Development, Western Australia. (2022), *Phalaris aquatica - environmental weed risk assessment 2022*. Department of Primary Industries and Regional Development, Western Australia, Perth. Report.

https://library.dpird.wa.gov.au/bs_wra/79

This report is brought to you for free and open access by the Biosecurity, pests, weeds and diseases research at Digital Library. It has been accepted for inclusion in Environmental Weed Risk Assessments by an authorized administrator of Digital Library. For more information, please contact library@dpird.wa.gov.au.



Department of Primary Industries and Regional Development

Environmental weed risk assessment

Phalaris (Phalaris aquatica)

Phalaris is a winter-active, temperate perennial grass native to the Mediterranean region which was introduced into Australia in the late 1800s. It is widely grown in pastures in south-eastern Australia but is a minor pasture species in high rainfall south-western Australia. Phalaris requires annual rainfall greater than 500mm and a growing season longer than 6 months to persist (Sanford 2006).

Phalaris has become naturalised and is common along roadsides in high rainfall areas of south-western Australia, particularly on medium-textured soils formed on fresh rock. Phalaris is a cool season grass and is unsuited to northern Western Australia (WA) even under irrigation.

Weed lists

National-international:

- Not listed in Weeds of Australia (398 weed species) <u>https://weeds.org.au/weeds-profiles/</u>
- "Phalaris is widely naturalised in southern and eastern Australia (i.e. in eastern New South Wales, the ACT, Victoria, Tasmania, south-eastern South Australia and south-western Western Australia).

....regarded as an environmental weed in Victoria, South Australia and New South Wales. It was also recently listed as a priority environmental weed in three Natural Resource Management regions. This species has been widely cultivated as a pasture grass in the temperate regions of Australia, particularly on the New South Wales tablelands. It is a weed of pastures, grasslands, open woodlands, roadsides, waste areas, disturbed sites, creek banks, riparian vegetation, floodplains and wetlands in the temperate and cooler sub-tropical regions of Australia" Weeds of Australia website Fact sheet Index (lucidcentral.org)

• In the Global Compendium of Weeds, phalaris is listed as an agricultural weed, casual alien, cultivation escape, environmental weed, naturalised, noxious weed, weed (Randall 2017).

Western Australia:

- "...A pasture grass, it occurs in pastures, drains, verges and disturbed sites from Moora to Narembeen, Albany and Manjimup..." (Hussey et al. 2007).
- Recorded as naturalised in the following IBRA Regions of WA: Geraldton sandplains, Avon wheatbelt, Swan coastal plain, Jarrah forest, Mallee and Warren (Keighery and Longman 2004).
- Naturalised taxa from nature reserves in WA (Keighery 1991).



Figure 1 Distribution of phalaris (*Phalaris aquatica*) in Australia (Source: 'The Australasian Virtual Herbarium')

Environmental weed risk assessment

Assessed using the 'Environmental weed risk assessment protocol for growing non-indigenous plants in the Western Australian rangelands' (Moore et al. 2022)

Region	Filter A	Filter B	
	Is the species a weed in similar environments in Australia or overseas?	Is the species likely to persist in the environment without management*?	Weed Risk Assessment rating
Kimberley	No	No	Negligible to low
Pilbara	No	No	Negligible to low
Gascoyne – Goldfields	No	No	Negligible to low
Agricultural area	Yes	Yes (>500mm AAR)	TBD

*Without management means no fertiliser, Rhizobia, irrigation, grazing management or control of competition from other species

References

Hussey BMJ, Keighery GJ, Dodd J, Lloyd SG, Cousens RD (2007) 'Western weeds. A guide to the weeds of Western Australia'. Second Edition. The Weeds Society of Western Australia Inc.

Keighery GJ (1991) Environmental weeds of Western Australia. Kowari, 2: 180-188.

- Keighery G, Longman V (2004) The naturalized vascular plants of Western Australia 1: Checklist, environmental weeds and distribution in IBRA regions. *Plant Protection Quarterly*, **19(1):** 12-32.
- Moore G, Munday C, Barua P (2022) 'Environmental weed risk assessment protocol for growing nonindigenous plants in the Western Australian rangelands', Department of Primary Industries and Regional Development, *Bulletin no. 4924*, Perth.

Randall RP (2017) 'Global compendium of weeds' (No. Ed. 3).

Sanford P (2006) Chapter 4, Temperate grasses. *In* 'Perennial pastures for Western Australia'. (Ed. GA Moore, P Sanford and T Wiley). Department of Primary Industries and Regional Development, Western Australia, Perth. Bulletin 4690.

Weeds of Australia database

https://keyserver.lucidcentral.org/weeds/data/media/Html/trifolium_repens.htm Site accessed 30 November 2021

Assessment by G Moore and N Nazeri January 2022

Important disclaimer

The Chief Executive Officer of the Department of Primary Industries and Regional Development and the State of Western Australia accept no liability whatsoever by reason of negligence or otherwise arising from the use or release of this information or any part of it.

Copyright © Department of Primary Industries and Regional Development, 2022